OFFICE OF STATEWIDE HEALTH PLANNING AND DEVELOPMENT Illogical Diagnosis Relationships

V0901 LOW BIRTH WEIGHT STATUS VERSUS LOW BIRTH WEIGHTS

- New 1/1/02

Guideline:

Status codes are to be used for patients who no longer have the initial condition, but may continue to manifest the consequences of it. Coding Clinic instructs that if a patient has a condition classifiable to the perinatal chapter (codes 760-779), it should be coded and these status V codes should not be used. Both the low birth weight status and low birth weights should not be used together.

V0901 Ex	xclusive check (if match, error) – Y438
Diagnosis Table	V21.3x	Low birth weight status
Relational Table	764.xx	Slow fetal growth and fetal malnutrition
V0901 Ex	xclusive check (if match, error) – Y439
Diagnosis Table	V21.3x	Low birth weight status
Relational Table	765.xx	Disorders relating to short gestation and low birth weight

41.

References: Coding Clinic for ICD-9-CM, AHA, 4th Quarter 2000, page 50.

ICD-9-CM Codebook, Tabular Section, Title Names and Excludes Notes under categories and codes

495 2003

OFFICE OF STATEWIDE HEALTH PLANNING AND DEVELOPMENT Illogical Diagnosis Relationships

V0902 POST-TERM PREGNANCY (40-42 weeks) VERSUS PROLONGED PREGNANCY

(greater than 42 weeks) – New 1-1-02

Guideline: Women who are between 40-42 weeks of pregnancy are considered post-term, if diagnosed

by the physician. Women who are past 42 weeks of pregnancy are considered prolonged pregnancy, if diagnosed by the physician. It is illogical to report codes for both the post-

term pregnancy and prolonged pregnancy.

V0902 Exclusive check (if match, error) – O017

Diagnosis Table 645.1x Post-term pregnancy (40-42 weeks)

Relational Table 645.2x Prolonged pregnancy (greater than 42 weeks)

<u>References:</u> Coding Clinic for ICD-9-CM, AHA, 4th Quarter 2000, page 43.

STOP!!!

496 2003